

Antarctic Circumpolar Current's Role in the Antarctic ice system: an overview

Douglas G. Martinson

Division of Ocean and Climate Physics,

Lamont-Doherty Earth Observatory

and

Department of Earth and Environmental Sciences Columbia University,

61 Route 9W, Palisades, NY 10964

dgm@ldeo.columbia.edu

1 845 365-8830

1 845 365-8736

1 Abstract

2 The Antarctic Circumpolar Current (ACC) provides fundamental control on the Antarctic ice
3 system. The tilt of the isopycnals of the ACC, in response to strong westerlies, serves to
4 thermally isolate the Antarctic continent from directly receiving the overwhelming subtropical
5 ocean surface heat. This same tilt provides the northern boundary of the polar seas; as such it
6 "contains" the statically stable cold fresh surface polar waters required for sea ice formation. In
7 this manner it effectively sets the northern limit for seasonal sea ice formation. The isopycnal tilt
8 also allows warm deep water to upwell to the surface near the continental margin in western
9 Antarctica where the ACC skirts the continental shelf, leading to excessive ocean heat flux to the
10 atmosphere in winter, and providing heat to melt the underside of the glacial ice.

11

12

13

14

15 **Keywords:** Antarctic Circumpolar Current, Ocean heat, Antarctica, glacial ice, sea level rise

16 **1.0 Introduction**

17 Currently the Antarctic Peninsula is undergoing extraordinary climate change¹. Investigation
18 of the role of ocean heat in this change illuminates a number of fundamental insights regarding
19 the important role of the Antarctic Circumpolar Current (ACC) in this and all other aspects of the
20 Antarctic ice system. In addition to substantial physical changes (e.g., Earth's most extreme
21 winter warming, and extensive sea ice and glacial ice loss), the western Antarctic Peninsula
22 (WAP) marine ecosystem is changing from that of a continental polar environment to one of
23 subpolar marine (Ducklow et al., 2007). The marine ecosystem is responding directly to changes
24 in the ice cover and ocean physical conditions, themselves responding to climate forcing; this
25 change is seen throughout the entire foodchain along the WAP (Montes-Hugo et al., 2009).

26 This paper describes the various roles played in this system by the ACC, using the extensive
27 (18 years in 2010) gridded data set of the Palmer Long Term Ecological Research project (Pal
28 LTER; Smith et al. 1995), supplemented by other available local data, including World Ocean
29 Circulation Experiment (WOCE) lines. ACC interactions within the continental shelf region as
30 revealed by LTER data along the WAP are used as a case study. WOCE lines help extend
31 relevant insights around Antarctica. As such, the paper presents an overview of how the ACC
32 influences the Antarctic cryosphere from existing studies.

33 **2.0 Background**

34 *2.1 Dramatic WAP Climate Change*

35 Dramatic climate change is being experienced in the present day WAP. Specifically: (1)
36 Vaughan et al. (2003) show that this region is undergoing the fastest warming on Earth in winter
37 (a time when the sun is near the horizon, contributing little radiative warming, and warm air cells

¹ A rather formal nomenclature exists whereby "climate change" refers to anthropogenic change, and "climate variability" refers to natural climate variability. In this document, I am not adhering to that nomenclature — climate change simply implies that the climate is changing, independent of attribution.

38 are uncommon), (2) Cook et al. (2005) show that 87% of the marine glaciers in the WAP are in
39 retreat, (3) Stammerjohn et al. (2008) show that all but a small area of the WAP perennial sea ice
40 disappeared between 1979 and 1999, with the winter seasonal sea ice season shortened by 3-4
41 months (>40% reduction), this decline continued through the most recent years including 2010
42 (Stammerjohn et al., 2010, and pers. com. for year 2010), (4) Meredith and King (2005) show
43 summer temperatures of the near-surface ocean rose more than 1 °C and salinities increased at
44 the same time as the atmospheric warming, and (5) Ducklow et al., 2007 and W. Fraser (personal
45 communication) show that the Adelie penguin colonies near Palmer Station in the WAP are
46 going extinct since the Adelies cannot adapt to changes in winter ice coverage¹.

47 The only substantial source of heat in the WAP winter is the ocean. The warmest water,
48 characterized by temperature (T_{\max}) and nutrient maxima, is Upper Circumpolar Deep Water
49 (UCDW) at 3.5 - 4°C above freezing in the ACC at the WAP. Martinson et al. (2008; hereafter
50 MSISV08), using the first 12 years of Pal LTER data, show that on the WAP continental shelf,
51 the heat content of the sub-surface water column (dominated by UCDW) has increased. During
52 the 1990s increased upwelling of UCDW onto the shelf explains ~84% of this increase; ~21% is
53 attributed to warmer UCDW².

54 While the WAP appears to be responding to global warming, the region also shows the
55 largest surface response to ENSO events outside of the tropics (e.g., Yuan and Martinson, 2001
56 and Yuan, 2004), and has been well documented to respond to changes in the strength of the
57 Southern Annular Mode (SAM). A positive bias in SAM leads to strengthening of the
58 atmospheric polar jet, driving more and stronger polar lows into the WAP, forcing upwelling
59 events (Marshall et al., 2004; Yuan, 2004). A similar response is realized from La Niña (Yuan,

¹ For a detailed popular description of this, see: Montaigne, F., 2010. "Fraser's Penguins", Henry Holt Press, 288 pp.

² Upwelling (84%) and warming (21%) of T_{\max} are independent, hence the total exceeds 100%.

60 2004). El Niño and negative SAM conditions each lead to the opposite (a reduction in storms
61 and upwelling events).

62 2.2 Data

63 Most of the data for this study have been acquired in the Pal LTER marine sampling grid
64 shown in Figure 1. The LTER project has collected shipboard data since 1991, including each
65 January since 1993 (i.e., summer snapshots). This has been recently supplemented by the
66 deployment of 5 Lamont temperature-string moorings¹ on the WAP continental shelf (mooring
67 strings contain temperature and pressure sensors at fixed locations from just above the seafloor to
68 a depth near 75 m depth, assuring sampling of all sub-surface water), and austral summer
69 launches of SLOCUM ocean gliders (1 in austral summer 2007; 2 in 2008; we now have a pool
70 of 3 such gliders in the project operated by Oscar Schofield of Rutgers). It is clear from the
71 research articles appearing in the 2008 Deep Sea Research Part II special issue "Palmer,
72 Antarctica Long Term Ecological Research", that the delivery of warm deep water to the
73 continental shelf of the WAP by the ACC plays a critical role in the physical and ecological
74 system. Hence data from this site are useful for revealing that aspect of the ACC's role in the
75 Antarctic ice system.
76

¹ Moorings designed by Lamont oceanographer Bruce Huber, designed to provide a temperature profile through the water column using fixed-depth sensors.




Figure 1: A. Regional map showing location of Pal LTER nominal ocean grid from which data for this study have been acquired. B. Ocean grid shows standard sample locations (solid squares) and geographic features labeled (top-to-bottom): An = Anvers Island (home of Palmer Station indicated by red cross), R = Renaud Island, L = Lavosier Island, Ad = Adelaide Island and MB = Marguerite Bay. 300.100 mooring, discussed in text, is green dot in grid center (number in center of mooring circles indicates year mooring was deployed). C. Temperature-Salinity plot shows collection of all summer profiles collected in grid since start of LTER project, identifying primary water masses, including UCDW supplying the ocean heat to the grid.

77 2.3 Physical Setting

78 The grid overlays the broad continental shelf of the WAP which is ~450 meters in depth
 79 (excluding canyons), running ~200 km in cross-shelf width and ~400 km along the WAP (the
 80 grid has been recently extended another ~300 km to the southwest). The southern boundary of
 81 the ACC, is most easily defined by the southern-most presence of UCDW (Orsi et al., 1995),
 82 given that the Southern ACC front is not always present. The southern boundary migrates to the

83 continental slope once it passes the Ross gyre and stays there throughout the entire SE Pacific
84 region (Orsi et al., 1995; MSISV08) along the continental rim of the Amundsen-Bellingshausen
85 Seas, riding up to the shelf-slope break in the WAP. This makes warm UCDW directly available
86 to the WAP shelf and the Amundsen Sea Embayment (where the major ice streams draining the
87 massive West Antarctic Ice Sheet, WAIS, enter the sea), allowing for easy ventilation to the
88 atmosphere and glacial melt. In this important respect, the West Antarctic continental shelf is
89 unique in Antarctica for this proximity of the ACC and delivery of warm UCDW¹ (Figure 2).




Figure 2: Polar projection shows proximity of Antarctic Circumpolar Current to entire Antarctic continent. ACC transports warm UCDW. Amundsen-Bellinghousen Seas are unique in entire Antarctic since here the ACC southern boundary follows the continental slope (up to shelf-slope break in WAP), delivering the warm UCDW directly to the shelf for ventilation of

¹ The ACC does briefly encounter the continental slope in short segments in the Indian Ocean sector, but it is not clear that that results in any significant delivery of UCDW to the continental shelf in those locations and if so, what the consequence is.

heat to the atmosphere and for access to the underside of glacial ice. Blue arrow head shows location of Amundsen Sea Embayment where major ice streams draining the WAIS flow into the sea. ACC is defined by those limiting geopotential surfaces that pass through Drake Passage (ACC based on climatological dynamic topography of Orsi et al., 1995); blue box shows location of LTER sampling grid. Blue-colored, labeled (with WOCE cruise track) lines cutting across ACC in 3 locations are WOCE diagrams identifying location of sections in Figure 3.

90 **3.0 Role of the ACC**

91 *3.1 Thermal Isolation*

92 First and foremost in the Antarctic thermal balance is the fact that the ocean water contains
93 >4000 times more thermal energy than air of the same volume (V) and temperature above
94 freezing (ΔT), reflecting the fact that thermal energy is a function of density and specific heat
95 capacity. And, given that strong solar forcing is absent half of the year, the ocean is the dominant
96 source of heat in winter. Thus, the presence of warm water near the Antarctic continent can
97 greatly moderate the atmospheric temperature and melt ice of any kind.

98 The most obvious role of the ACC is the fact that it thermally isolates Antarctica, as
99 suggested by Kennett (1977), by keeping warm subtropical surface waters from approaching the
100 continent. In particular, the lack of physical boundaries along its path prevents development of
101 zonal pressure gradients driving meridional flow. Also, the westerlies force the ACC so that the
102 stratified water column is tilted higher in the south. Together, these prevent the warm subtropical
103 surface waters north of the ACC from directly moving south across the northern edge of the
104 ACC, as seen in the WOCE temperature sections of Figure 3. Each section from the SE Pacific
105 (P16), mid-Atlantic (S2) and Indian Ocean (I6), clearly shows that the tilt prevents the warm
106 subtropical surface waters, containing an immense heat content, from moving south directly to
107 Antarctica, helping to keep it and its surrounding oceans cold (establishing Kennett's "steep
108 surface water temperature gradient between polar and tropical regions"). Note that this is not the
109 case in the North Atlantic where warm subtropical waters (in the North Atlantic Drift) are

110 transported directly to the Nordic Seas impeding sea ice extension to the south (Untersteiner,
111 1988) near Fram Strait.




Figure 3: 3-color WOCE Temperature sections along the tracks crossing the ACC in polar projection (in Figure 2) labeled same as those sections. Sections show that tilt of the isopycnals associated with ACC prevent warm tropical waters (red) from reaching Antarctic continental margin. Warm UCDW layer (white) slips along the tilted isopycnals to reach the continental slope in section P19 along the WAP, and elsewhere only reaches the northern limit of the polar waters.

112 While the ACC delimits the southern boundary of subtropical surface waters, it also
113 "contains" the cold fresh surface polar waters required for sea ice formation, so it effectively sets
114 the northern limit for seasonal sea ice formation. That limit defines the extent of the Antarctic
115 polar oceans (as seen in the spatial distribution of Winter Water, generated by the deep winter
116 mixed layers resulting from destabilizing brine rejection associated with sea ice growth, Toole,
117 1981). In some regions the weak stratification of the Southern Ocean water column leads to a

118 strong bathymetric control on the location of the ACC. Paleoceanographic studies indicating a
119 northward migration of the sea ice fields during ice age climates (e.g., CLIMAP reconstruction,
120 1981 and Gersonde et al., 2005) are also revealing a northward migration of the ACC. In those
121 regions showing strong bathymetric control on the ACC, this northward migration implies
122 changes releasing the ACC from the bathymetric control.

123 *3.2 Modern heating of WAP waters over the shelf*

124 While the ACC inhibits warm subtropical surface waters from southward extension into the
125 polar oceans, the warm deep water (as UCDW in the ACC) is elevated from depth along the
126 tilted isopycnal surfaces. In the WAP, the warm deep water is elevated to the depth of the floor
127 of the continental shelf (section P19 in Figure 2) allowing some of this warm water direct access
128 to glacier margins. In the other ocean basins the warm water enters the polar gyres, providing the
129 heat that dominates the ocean stability and limits the thickness of the seasonal sea ice
130 (Martinson, 1990, Martinson and Iannuzzi, 1998, 2003).

131 MSISV08 document the ocean heat content on the shelf, and show that when including
132 historical data in the same vicinity of the LTER sampling grid there has been an exponential
133 increase in this heat content over the decades since the 1960s. Fundamental to this increased
134 ocean heat content is the applicability of the WAP findings to the Amundsen Sea Embayment.
135 Recent research there indicates that indeed it is the (same) warmed UCDW water in that region
136 responsible for the accelerated glacial melt there (e.g., Shepherd et al., 2002, Payne et al., 2004).
137 This, as well as the robustness of the temporal increase in ocean heat is the focus of current
138 research and will be presented in a separate publication.

139 **4.0 Discussion and Conclusions**

140 *4.1 Discussion.*

141 The ACC is playing a crucial role in the Antarctic ice system. It serves to minimize the flux
142 of surface subtropical heat into the polar oceans, thus thermally isolating the Antarctic continent.
143 For that same reason, it also serves as a northern boundary to the polar seas, containing the fresh
144 cold stable surface waters necessary for sea ice formation and limiting the northern extent of the
145 seasonal sea ice fields. Some of the modern ACC path is controlled by bathymetry;
146 paleoceanographic data indicating north-south shifts in the sea ice fields suggest a release from
147 that bathymetric control — presumably something that could be achieved if, for example,
148 enough sea ice melted at the northern edge of the polar gyres, forming a strongly stratified
149 surface layer, limiting penetration of surface forcing and consequently decreasing the
150 bathymetric control.

151 The ACC transports warm water (relative to freezing), in the form of UCDW, around the
152 continent. The edge of the East Antarctic ice cap is buffered from this heat by large polar gyres
153 (the Weddell and Ross gyres) and by a seafloor bathymetry that steers it through the Indian
154 Ocean minimizing its contact with the continental shelf in that region. Only after it passes the
155 Ross Sea gyre does it immediately move continent-wise likely owing to the wavenumber 3
156 atmospheric circulation pattern (van Loon and Jenne, 1972), where it continues its journey
157 skirting the edge of the continental shelf making the heat easily accessible to the region most
158 susceptible to draining the WAIS and WAP glaciers. In the WAP it is at a depth where a simple
159 offshore surface flow (MSISV08) will draw the warmest water onto the shelf.

160 *4.2 Conclusions.*

161 It is shown that the tilt of the isopycnals of the ACC contributes to the thermal isolation of
162 Antarctica, and in a complementary sense, serves to isolate the seasonal sea ice fields. That same
163 tilt delivers warm deep water to the continental shelves of the western Antarctica providing heat

164 to a region sensitive to glacial melt. Still at question is the degree to which the warm deep water
165 directly contributes to the accelerated glacial melt currently being experienced by the WAIS via
166 the ice streams (Pine Island, Thwaites and Smith) flowing into the Amundsen Sea Embayment.

167 **Acknowledgments**

168 Thanks to the organizers of the ACE Symposium for asking me to participate in that
169 interesting forum, and providing helpful comments for this paper. And, thanks, as always to
170 Richard A. Iannuzzi who contributed heavily to the analyses and figures, and provided insightful
171 science and editorial suggestions. Deepa Manandhar contributed to the figures and references.
172 Thanks to the crew, staff and Raytheon Polar Services and our own LTER science support teams
173 on our numerous LTER cruises aboard the RVIB LM Gould; all cooperated fully to help us
174 acquire the data needed to answer the questions we were addressing. This work was supported by
175 NSF grants: LTER 0217282 and 0823101, and SASSI. NOAA grant: NA08OAR4320912. This
176 paper is LTER contribution xxx and Lamont contribution xxx.

177

178 **References**

179 AchutaRao, K.M., Ishii, M., Santer, B.D., Gleckler, P.J., Taylor, K.E., Barnett, T.P., Pierce,
180 D.W. Stouffer, R.J., Wigley, T.M.L., 2007. Simulated and observed variability in ocean
181 temperature and heat content. *Proceedings of the National Academy of Sciences* 104 (26),
182 10768-73.
183 Alley, R.B., Clark, P.U., Huybrechts, P., Joughin, I., 2005. Ice-sheet and sea-level changes.
184 *Science* 310, 456-460.
185 Anderson, J.B., 2002. *Antarctic Marine Geology*, Cambridge University Press, Cambridge, UK.

186 Barnett, T.P., Pierce, D.W., Shnur, R., 2001. Detection of anthropogenic climate change in the
187 world's oceans. *Science* 292, 270-274.

188 Barnett, T.P., Pierce, D.W., AchutaRao, K.M., Gleckler, P.J., Santer, B.D., Gregory J.M.,
189 Washington, W.M. 2005. Penetration of human –induced warming into the world’s oceans.
190 *Science* 309, 284-287.

191 Böning, C.W., Dispert, A., Visbeck, M., Rintoul S.R., Schwartzkopf, F.U., 2008. The response
192 of the Antarctic Circumpolar Current to recent climate change. *Nature Geoscience* 1, 864-869.

193 Cook, A.J., Fox, A.J., Vaughan, D.G., Ferrigno, J.C., 2005. Retreating glacial fronts on the
194 Antarctic Peninsula over the past half-century. *Science* 308, 541-544.

195 Ducklow, H.W., Baker, K.S., Martinson, D.G., Quetin, L.B., Ross, R.M., Smith, R.C.,
196 Stammerjohn, S.E., Vernet, M., Fraser, W., 2007. Marine pelagic ecosystems: The West
197 Antarctic Peninsula. *Philosophical Transactions of the Royal Society of London Special Theme*
198 *Issue, Antarctic ecology: From genes to ecosystems* 362, 67-94.

199 Gersonde, R., Crosta, X., Abelmann, A., Armand, L., 2005. Sea-surface temperature and sea ice
200 distribution of the Southern Ocean at the EPILOG Last Glacial Maximum a circum-Antarctic
201 view based on siliceous microfossil records, *Quaternary science reviews*, 24, 869-896.
202 doi:[10.1016/j.quascirev.2004.07.015](https://doi.org/10.1016/j.quascirev.2004.07.015)

203 Gille, S.T., 2002. Warming of the Southern Ocean since the 1950s. *Science* 295, 1275-1277.

204 Gille, S.T., 2008. Decadal-scale temperature trends in the southern hemisphere ocean. *Journal of*
205 *Climate* 21, doi: 10.1175/2008JCLI2131.1.

206 Jacobs , S.S., Hellmer, H.H. and Jenkins, A., 1996. Antarctic ice sheet melting in the southeast
207 Pacific. *Geophysical Research Letters* 23 (9), 957-960.

208 Kennett, J.P., 1977. Cenozoic evolution of Antarctic glaciation, the circum-Antarctic ocean, and
209 their impact on global paleoceanography. *Journal of Geophysical Research* 82 (27), 3843-3860.

210 Levitus, S., Antonov, J., Boyer T.P., Stephens, C., 2000. Warming of the world ocean. *Science*
211 287, 2225-2229.

212 Levitus, S., Antonov, J.I., Wang, J., Delworth, T.L., Dixon, K.W., Broccoli, A.J., 2001.
213 Anthropogenic warming of earth's climate system. *Science* 292, 267-270.

214 Levitus, S., Antonov, J.I., Boyer, T.P., 2005. Warming of the world ocean, 1955-2003.
215 *Geophysical Research Letters* 32, doi: 10.1029/2004GL021592.

216 Marshall G.J, Stott P.A., Turner J., Connolley W.M., King J.C., Lachlan-Cope T.A., 2004.
217 Causes of exceptional atmospheric circulation changes in the southern hemisphere. *Geophysical*
218 *Research Letters* 31, L14205, doi:101029/2004GL019952.

219 Martinson, D.G., 1990. Evolution of the Southern Ocean winter mixed layer and sea ice: open
220 ocean deepwater formation and ventilation. *Journal of Geophysical Research* 95, 11641-11654.

221 Martinson, D.G., Iannuzzi, R.A., 1998. Antarctic ocean-ice interaction: Implications from ocean
222 bulk property distributions in the Weddell Gyre, in: Jeffries, M. [Ed.], *Antarctic Sea Ice:*
223 *Physical Processes, Interactions and Variability*. Antarctic Res. Series, vol. 74, American
224 Geophysical Union, Boston, pp. 243-71.

225 Martinson, D.G., Iannuzzi, R.A., 2003. Spatial/temporal patterns in Weddell Gyre characteristics
226 and their relationship to global climate. *Journal of Geophysical Research* 108 (C4)
227 doi:10.1029/2005GL024042.

228 Martinson, D.G., Stammerjohn, S.E., Iannuzzi, R.A., Smith R.C., Vernet, M., 2008. Western
229 Antarctic Peninsula physical oceanography and spatio-temporal variability. *Deep-Sea Research*
230 II 55, 1964-1987

231 Meredith, M.P., King, J.C., 2005. Rapid climate change in the ocean to the west of the Antarctic
232 Peninsula during the second half of the twentieth century, *Geophysical Research Letters*, 32,
233 L19604. doi:10.1029/2005GL024042.

234 Moffat, C., Owens, B., Beardsley, R.C., 2009. On the characteristics of circumpolar deep water
235 intrusions to the west Antarctic Peninsula continental shelf. *Journal of Geophysical Research*,
236 114, C05017, doi:10.1029/2008JC004955.

237 Montes-Hugo, M.A., Doney, S.C., Ducklow, H.W., Fraser, W., Martinson, D., Stammerjohn,
238 S.E., Schofield, O., 2009. Recent changes in phytoplankton communities associated with rapid
239 regional climate change along the Western Antarctic Peninsula. *Science* 323, 1470-1473.

240 Orsi, A.H., Whitworth, T., Nowlin, W.D., 1995. On the meridional extent and fronts of the
241 Antarctic Circumpolar Current. *Deep Sea Research I*, 42 (5), 641-673.

242 Payne, A.J., Vieli, A., Shepherd, A.P., Wingham, D.J., Rignot, E., 2004. Recent dramatic
243 thinning of largest West Antarctic ice stream triggered by oceans. *Geophysical Research Letters*
244 31, L23401, doi:10.1029/2004GL021284.

245 Schlitzer, R., 2009. Ocean Data View, <http://odv.awi.de>.

246 Shepherd, A., Wingham, D. and Mansley, J., 2002. Inland thinning of the Amundsen Sea sector,
247 west Antarctica, *Geophysical Research Letters* 29(10), 2-1.

248 Smith, R.C., Baker, K.S., Fraser, W.R., Hofmann, E.E., Karl, D.M., Klinck, J.M., Quetin, L.B.,
249 Prezelin, B.B., Ross, R.M., Trivelpiece, W.Z., Vernet, M., 1995. The Palmer LTER: A long-term
250 ecological research program at Palmer Station, Antarctica. *Oceanography* 8 (3), 77-86.

251 Stammerjohn, S.E., Martinson, D.G., Smith, R.C., Iannuzzi, R.A., 2008. Sea ice in the western
252 Antarctic Peninsula region: Spatio-temporal variability from ecological and climate change
253 perspectives. *Deep Sea Research Part II*, 55 (18-19), 2041-2058, doi:10.1016/j.dsr2.2008.04.026.

254 Steig, E.J., Schneider, D.P., Rutherford, S.D., Mann, M.E., Comiso, J.C., Shindell, D.T., 2009.
255 Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year.
256 Nature 457 (22), doi:10.1038 /nature 7669.

257 Thoma, M., Jenkins, A., Holland, D., Jacobs, S., 2008. Modelling Circumpolar Deep Water
258 intrusions on the Amundsen Sea continental shelf, Antarctica. Geophysical Research Letters, 35,
259 L18602, doi:10.1029/2008GL034939.

260 Toole, J.M., 1981. Sea ice, winter convection, and the temperature minimum layer in the
261 Southern Ocean, Journal of Geophysical Research, 86, 8037-1047.

262 Untersteiner, N., 1988. On the ice and heat balance in Fram Strait. Journal of Geophysical
263 Research 93 (C1), 527-531.

264 van Loon, H., Jenne, R.L., 1972. The zonal harmonic standing waves in the southern
265 hemisphere. Journal of Geophysical Research 77 (6), 992-1003.

266 Vaughan, D.G., Marshall, G.J., Connolley, W.M., Parkinson, C., Mulvaney, R., Hodgson, D.A.,
267 King, J.C., Pudsey, C.J., Turner, J., 2003. Recent rapid regional climate warming on the
268 Antarctic Peninsula. Climatic Change 60, 243-274.

269 Yuan, X., Martinson, D.G., 2001. The Antarctic Dipole and its predictability. Geophysical
270 Research Letters 28 (18), 3609-3612.

271 Yuan, X., 2004. ENSO-related impacts on Antarctic sea ice: a synthesis of phenomenon and
272 mechanisms. Antarctic Science 16 (4), 415-425.